



PERIOD / EPOCH		CONODONT ZONES
ORDOVICIAN	CARADOC	
	LLANVIRN	<i>Phragmodus-Plectodina</i> <i>Histiodelia holodentata</i>
		<i>Histiodelia altitrons</i>
	ARENIG	<i>Jumudontus gananda</i> <i>Oepikodus communis</i> <i>Prioniodus elegans</i> – <i>Bergstroemognathus extensus</i> <i>Drepanolostodus</i>
	TREMADOC	

CONODONT ALTERATION INDEX
1
1 – 1.5
1.5
2
2 – 3
3

Note: Where an age is 'undifferentiated zone x to zone y', both zone colours are shown in an expanded column width

- Conglomerate
- Pebbly sandstone
- Sandstone
- Siltstone
- Shale, claystone
- Sandy limestone
- Sandy dolostone
- Argillaceous or silty limestone
- Argillaceous or silty dolostone
- Limestone
- Dolostone
- Calcareous
- Dolomitic
- Limestone or dolostone interbeds
- Halite
- Anhydrite or gypsum
- Glauconitic
- Granitic basement
- Metamorphic basement
- Unconformity
- Interpreted correlation of formal geological unit
- Tentative correlation of formal geological unit
- Tentative correlation of informal geological unit
- ?Ash bed
- Granitic pebbles – clasts

Compiled by P.W. Haines
Edited by M. Aphorpe
Cartography by L. E. Fogarty
Published by the Geological Survey of Western Australia. Digital and hard copies of this plate are available from the Information Centre, Department of Industry and Resources, 100 Plain Street, East Perth, WA, 6004. Phone (08) 9222 3459, Fax (08) 9222 3444, Web www.doir.wa.gov.au/gswa, Email geological_survey@doir.wa.gov.au
The recommended reference for this plate is:
HAINES, P.W., 2004, Stratigraphic correlations, Munro 1 to Hedonia 1 (Willara Sub-basin to Broome Platform), in Depositional facies and regional correlations of the Ordovician Goldwyer and Nita Formations, Canning Basin, Western Australia, with implications for petroleum exploration by P.W. HAINES: Western Australia Geological Survey, Record 2004/7, Plate 1.